

Color- $SU(3)$ -Ginzburg-Landau Effective Potential for Order Parameter with 3×3 Symmetry

E. Nakano, T. Suzuki and H. Yabu

*Department of Physics, Tokyo Metropolitan University, 1-1 Minami-Ohsawa, Hachioji, Tokyo
192-0397, Japan*

(February 7, 2008)

Abstract

Ginzburg-Landau effective potential is studied for the order parameter that transforms in the $(3, 3)$ representation under the color $SU(3)$ group. All the $SU(3)$ invariant terms to the fourth-order of the order parameter are classified and the effective potential is constructed in its most general form. The conditions which stabilize the condensed phases are also obtained. As applications, the classification of the condensed phase is discussed in some special cases and the Higgs phenomenon associated with symmetry breaking is studied by introducing a coupling to the gauge boson.

I. INTRODUCTION

Recently, quark-pair condensation phenomena (color-superconductor) have received much interest in the high-density nuclear physics [1,2].

In a high-density quark matter, quarks make a correlated Cooper pair due to the attractive one-gluon-exchange [3,6] or instanton-induced interactions [4]. A lot of studies have been done for physical properties of color superconductors on the basis of the QCD-inspired models [5–9].

From the symmetry point of view, the color-superconductor is characterized by the spontaneous symmetry breaking of the color $SU_C(3)$ symmetry. Let's consider the quark field operator $q_\alpha(r)$ with the color index $\alpha = 1, 2, 3$ (other degrees of freedom, flavors and spins, are represented, e.g., as $q = u_\uparrow$). They transform as a vector (the **3**-representation) under the $SU_C(3)$ transformation. (Irreducible representations of $SU(3)$ are denoted by their dimensions, e.g. **1**, **3**, **3***, etc.) The color-superconductor is a long-range ordered state, the order parameter of which is a two-body one: $\Psi_{\alpha,\beta}(r) = \langle q_\alpha(r)q'_\beta(r) \rangle$. It is a second-rank tensor under $SU_C(3)$ with nine components, which makes a reducible $\mathbf{3} \times \mathbf{3}$ -representation.

Different from the systems with a scalar order parameter (metal superconductor, superfluid ^4He , etc.), a variety of symmetry-breaking patterns shows up in the condensed states with the tensor order parameter. A typical example is the superfluid ^3He , where, due to the p-wave nature of the Cooper pair, the order parameter has a tensor-type spin structure, $\Phi_{i,j}$ ($i, j = 1, 2, 3$); it transforms as the $\mathbf{3} \times \mathbf{3}$ -representation in the orbital and spin rotations $G = SO_L(3) \otimes SO_S(3)$ (also with the gauge $U(1)$ symmetry). Corresponding to several symmetry breaking patterns, many different condensed phases show up in the superfluid ^3He : A-phase ($H = (S^2 \times SO(3))/Z_2$), B-phase ($R_B = S_1 \times SO_a(3)$), and so on. As discussed later, such a phase structure should exist also for the $SU(3)$ - $\mathbf{3} \times \mathbf{3}$ -type order parameter, and it is very interesting to study the structures for such a novel symmetry breaking case [7]. (It should be noted that the $\mathbf{3} \times \mathbf{3}$ representation of $SU(3)$ is a complexification of that of $SO(3)$, so that the color superconducting state can be considered as a kind of the group

extension for the superfluid ^3He .)

The systems mentioned above are well described by the thermodynamic effective free energy that is constructed as a function of the order parameter up to the fourth-order, i.e., Ginzburg-Landau method [10]. This method incorporates the coherence nature of the pair condensation and describes the second order phase transition. It was also applied to the color-superconductor in some special cases [11].

In this paper, we discuss phase structures of the system with $SU(3)$ - $\mathbf{3} \times \mathbf{3}$ order parameter based on the Ginzburg-Landau method: the general form of the Ginzburg-Landau effective potential is presented to the fourth-order of the order parameter $\Phi_{\alpha,\beta}$, and analyzed in detail. As applications, the classification of the condensed phase and Higgs phenomenon are discussed in some special cases.

II. STRUCTURE OF $SU(3)$ - $\mathbf{3} \times \mathbf{3}$ ORDER PARAMETER

Let's consider the system with the symmetry group $SU(3)$, where the order parameter is given by the 3-by-3 matrix : $\Psi = (\Psi_{\alpha,\beta})$. For $g \in SU(3)$, it transforms as

$$\Psi \longrightarrow g \Psi^t g, \quad (1)$$

where $^t g$ is a transposition of g . It makes a $\mathbf{3} \times \mathbf{3}$ -representation of $SU(3)$.

The system should also be invariant under the $U(1)$ gauge transformation:

$$\Psi \longrightarrow e^{i\phi} \Psi. \quad (2)$$

Because of the irreducible decomposition $\mathbf{3} \times \mathbf{3} = \mathbf{3}^* + \mathbf{6}$, the order parameter $\Psi = (\Psi_{\alpha,\beta})$ can be decomposed into the irreducible components: $\Psi = S + A$, where S and A are the symmetric and antisymmetric components:

$$S = \frac{1}{2}(\Psi + ^t\Psi), \quad A = \frac{1}{2}(\Psi - ^t\Psi). \quad (3)$$

Since the permutation symmetry for indices (α, β) are conserved under the $SU(3)$ -transformation, the S and A become basis for the subrepresentations of $SU(3)$. By dimension counting, we find that the S and A correspond to the $\mathbf{6}$ and $\mathbf{3}^*$ representations.

From this consideration, symmetry breaking patterns for the condensed phases are classified into four types:

$$1) \text{ No-condensate} \quad A = S = 0, \quad (4)$$

$$2) \text{ } \mathbf{3}^* \text{-condensate} \quad A \neq 0, \quad S = 0, \quad (5)$$

$$3) \text{ } \mathbf{6} \text{-condensate} \quad A = 0, \quad S \neq 0, \quad (6)$$

$$4) \text{ Mixed-condensate} \quad A \neq 0, \quad S \neq 0. \quad (7)$$

Because of the $SU(3)$ invariance of the system, for $g \in SU(3)$, order parameters (A, S) and $(gA^t g, gS^t g)$ represent the same condensed states, so that we can decompose the order parameters A and S ;

$$A = e^{i\theta} G A_0^t G, \quad S = e^{i\theta} G S_0^t G, \quad (8)$$

where $G \in SU(3)$ and $e^{i\theta} \in U(1)$ are the Goldstone degrees of freedom which can be eliminated by the $SU(3)$ and $U(1)$ gauge transformations. Remaining parts A_0 and S_0 in eq. (8) determine essentially scales of the symmetry breaking. As proved in appendix A, they can be parametrized as

$$A_0 = \begin{pmatrix} 0 & \phi_3 & -\phi_2 \\ -\phi_3 & 0 & \phi_1 \\ \phi_2 & -\phi_1 & 0 \end{pmatrix}, \quad S_0 = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix}, \quad (9)$$

where $d_{1,2,3}$ are real and $\phi_{1,2,3}$ are complex parameters: $\phi_i = |\phi_i| e^{i\theta_i}$. We take the parametrization in eqs. (8) and (9) as a standard form and call d_i and ϕ_i normal coordinates of the $\mathbf{3} \times \mathbf{3}$ order parameter.

III. EFFECTIVE POTENTIAL FOR $SU(3)$ - $\mathbf{3} \times \mathbf{3}$ ORDER PARAMETER

In the Ginzburg-Landau theory, the effective potential V_{eff} for the order parameter Ψ plays a central role. It should be a invariant function of Ψ for the $SU_C(3)$ and $U(1)$ gauge transformations, (1) and (2). For a minimal Ginzburg-Landau model, it is necessary to construct V_{eff} to the fourth-order of Ψ .

The lowest order contributions to V_{eff} are the second-order terms in Ψ and there exist such two terms constructed from A and S :

$$V_{\text{eff}}^{(2)} = a_1 \text{Tr} A^* A + a_2 \text{Tr} S^* S. \quad (10)$$

All other second-order terms can be shown to vanish or to be transformed into the above form (e.g., $\text{Tr} A^* S = 0$ due to the antisymmetry of $A^* S$.)

In a similar way, we can construct a general fourth-order potential $V_{\text{eff}}^{(4)}$:

$$\begin{aligned} V_{\text{eff}}^{(4)} = & b_1 (\text{Tr} A^* A)^2 + b_2 (\text{Tr} S^* S)^2 + b_3 \text{Tr} S^* S S^* S \\ & + b_4 \text{Tr} A^* A \text{Tr} S^* S + b_5 \text{Tr} A^* A S^* S + b_+ (\text{Tr} A^* S A^* S + \text{c.c.}) + i b_- (\text{Tr} A^* S A^* S - \text{c.c.}). \end{aligned} \quad (11)$$

In the same way as for $V^{(2)}$, any other fourth-order terms can be transformed into the above form: e.g., $(\text{Tr} A^* A)^2 = 2 \text{Tr} A^* A A^* A$.

Using eq. (8), the potentials $V_{\text{eff}}^{(2,4)}$, eqs. (10) and (11), can be rewritten with the normal coordinates $(d_i, |\phi_i|, \theta_i)$ defined in eq. (9). For simplification, we introduce the variables $R \equiv \sqrt{|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2}$, $D \equiv \sqrt{d_1^2 + d_2^2 + d_3^2}$ and the normalized coordinates: $r_i \equiv |\phi_i|/R$ and $e_i \equiv d_i/D$, then eqs. (10) and (11) become

$$V_{\text{eff}}^{(2+4)} = g_R R^2 + g'_R R^4 + g_D D^2 + (g'_D - h_D x) D^4 + (g''_D + h''_D u) R^2 D^2, \quad (12)$$

where the coefficients g_i are defined by linear combinations of a_i and b_i in eqs. (10) and (11):

$$\begin{aligned} g_R = -2a_1, \quad g'_R = 4b_1, \quad g_D = a_2, \quad g'_D = b_2 + b_3, \\ h_D = 2b_3, \quad g''_D = -2b_4 - b_5, \quad h''_D = b_5. \end{aligned} \quad (13)$$

The variables x and u in eq. (12) are defined by

$$x \equiv e_1^2 e_2^2 + e_2^2 e_3^2 + e_3^2 e_1^2, \quad u \equiv f_1 r_1^2 + f_2 r_2^2 + f_3 r_3^2, \quad (14)$$

where

$$\begin{aligned} f_1 = e_1^2 + c \sin(2\theta_1 + \delta) e_2 e_3, \quad f_2 = e_2^2 + c \sin(2\theta_2 + \delta) e_3 e_1, \quad f_3 = e_3^2 + c \sin(2\theta_3 + \delta) e_1 e_2. \end{aligned} \quad (15)$$

The constants c and δ in eq. (15) are obtained from the coefficients $b_{\pm}$ in eq. (11) as $c = -4\sqrt{b_+^2 + b_-^2}/b_5$ and $\tan \delta = b_+/b_-$.

IV. DOMAIN OF VARIABLES

Now we obtain the general form of the effective potential to the fourth-order, which is a function of the variables (R, D, x, u) . Let's consider the domain of these parameters.

First, the variables R and D are defined as the lengths of $|\phi_i|$ and d_i , so that $R > 0$ and $D > 0$.

We introduce new variables:

$$0 \leq p_i \equiv e_i^2 \leq 1, \quad 0 \leq q_i \equiv r_i^2 \leq 1, \quad -1 \leq \lambda_i \equiv \sin(\theta_i + \delta) \leq 1, \quad (i = 1, 2, 3) \quad (16)$$

which satisfies the constraints: $p_1 + p_2 + p_3 = 1$ and $q_1 + q_2 + q_3 = 1$. Geometrically, possible ranges in the (p_1, p_2, p_3) or (q_1, q_2, q_3) space become the two-dimensional region inside the equilateral triangle with the vertices $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$. Using these variables, the variables x and u are represented by

$$x = p_2 p_3 + p_3 p_1 + p_1 p_2 = \frac{1}{2} \left\{ 1 - (p_1^2 + p_2^2 + p_3^2) \right\}, \quad u = f_1 q_1 + f_2 q_2 + f_3 q_3, \quad (17)$$

where $f_1 = p_1 + c\lambda_1\sqrt{p_2 p_3}$.

Let's consider the domain of $x = p_1 p_2 + p_2 p_3 + p_3 p_1$. It is clear that $x \geq 0$ because $p_{1,2,3} \geq 0$. Using the inequalities $(p_i + p_j)^2 \geq 4p_i p_j$, we obtain

$$1 = (p_1 + p_2 + p_3)^2 = \frac{(p_1 + p_2)^2 + (p_2 + p_3)^2 + (p_3 + p_1)^2}{2} + x \geq \frac{4x}{2} + x = 3x. \quad (18)$$

Thus, the variable x takes the values $0 \leq x \leq 1/3$.

The domain of $u = f_1 q_1 + f_2 q_2 + f_3 q_3$ depends on the value of x . Here, we summarize the derivation:

1. *Minimum/maximum of u for fixed f_i .* Because $0 \leq q_i \leq 1$ with $q_1 + q_2 + q_3 = 1$, the domain of u becomes $\min\{f_1, f_2, f_3\} \leq u \leq \max\{f_1, f_2, f_3\}$.
2. *Minimum/maximum of $f_1 = p_1 + c\lambda_1\sqrt{p_2 p_3}$ for fixed p_i .* Without loss of generality, we can assume $c \geq 0$ because $-1 \leq \lambda_1 \leq 1$. For fixed p_i , the variable f_1 takes a minimum value $p_1 - c\sqrt{p_2 p_3}$ at $\lambda_1 = -1$ and a maximum value $p_1 + c\sqrt{p_2 p_3}$ at $\lambda_1 = 1$. Because

f_i are defined symmetrically for the index ($i = 1, 2, 3$), similar results are obtained for f_2 and f_3 .

3. *Range of u for fixed p_i .* Combining the above results 1. and 2. , we obtain the possible range of u for fixed p_i : $f_1^- \leq u \leq f_1^+$, where

$$f_1^- = p_1 - c\sqrt{p_2 p_3}, \quad f_1^+ = p_1 + c\sqrt{p_2 p_3}. \quad (19)$$

4. Finally the domain of u for the fixed value of x is obtained by $\min \{f_1^-, f_2^-, f_3^-\} \leq u \leq \max \{f_1^+, f_2^+, f_3^+\}$ where min and max are for $0 \leq p_i \leq 1$ with $p_1 + p_2 + p_3 = 1$.

Thus, the evaluation of the domain of u has been reduced to those of $\min(f_-)$ and $\max(f_+)$, which are somewhat complicated because they depend not only on the coefficient c but also on the variable x . The calculations for them are given in appendix B, and we show only the final results for the domain of (x, u) below.

The results are given for $(0 \leq c \leq 2)$ and $(2 \leq c)$, which are respectively shown in figure 1(a) and (b). These figures show upper and lower bounds of u for fixed x .

a : $0 \leq c \leq 2$

$$\text{Upper bound of } u : \frac{2-c}{3}\sqrt{1-3x} + \frac{1+c}{3} \quad (20)$$

$$\text{Lower bound of } u : \begin{cases} -c\sqrt{x} & (0 \leq x \leq \frac{1}{4}) \\ -\frac{2+c}{3}\sqrt{1-3x} + \frac{1-c}{3} & (\frac{1}{4} \leq x \leq \frac{1}{3}) \end{cases} \quad (21)$$

The segment $\overline{AB}$ in figure 1(a) corresponds to eq. (20), and $\overline{OC}$ and $\overline{CD}$ to two equations in (21).

b : $2 \leq c$

$$\text{Upper bound of } u : \begin{cases} -\frac{c-2}{3}\sqrt{1-3x} + \frac{1+c}{3} & (0 \leq x \leq \frac{(2c-1)^2}{4(c^2-c+1)^2}) \\ c\sqrt{x} & (\frac{(2c-1)^2}{4(c^2-c+1)^2} \leq x \leq \frac{1}{4}) \\ \frac{c-2}{3}\sqrt{1-3x} + \frac{1+c}{3} & (\frac{1}{4} \leq x \leq \frac{1}{3}) \end{cases} \quad (22)$$

$$\text{Lower bound of } u : \begin{cases} -c\sqrt{x} & (0 \leq x \leq \frac{1}{4}) \\ -\frac{2+c}{3}\sqrt{1-3x} + \frac{1-c}{3} & (\frac{1}{4} \leq x \leq \frac{1}{3}) \end{cases} \quad (23)$$

The segments $\overline{AE}, \overline{EF}$ and $\overline{FB}$ in figure 1(b) correspond to three divisions in eq. (22), and $\overline{OC}$ and $\overline{CD}$ to two equations in (23).

V. STABILITY CONDITION

For the system with the effective potential V_{eff} to be physically meaningful, it should be positive definite when the variables R or D take infinitely large value. (If not, any states with finite R and D become unstable.)

This stability (positive definite) condition gives constraints on the parameters g_i and h_i in eq. (12) and we derive them in this section. First, we consider special cases:

1. When $D = 0$ and $R \rightarrow \infty$, eq. (12) become $V_{\text{eff}} \sim g'_R R^4$
2. When $R = x = 0$ and $D \rightarrow \infty$, we obtain $V_{\text{eff}} \sim g'_D D^4$.

From the positive definiteness of V_{eff} in these limits, the first two constraints can be obtained:

$$g'_D \geq 0, \quad g'_R \geq 0. \quad (24)$$

Hereafter, we consider the cases of $g'_D > 0$ and $g'_R > 0$ only. (We can treat the cases that $g'_D = 0$ or $g'_R = 0$ with taking the limit of those in $g'_D > 0$ and $g'_R > 0$.)

We replace the variables R and D by the scaled ones: $R \rightarrow R/(g'_R)^{\frac{1}{4}}$ and $D \rightarrow D/(g'_D)^{\frac{1}{4}}$, then eq. (12) becomes

$$V_{\text{eff}} = -GR^2 + R^4 - FD^2 + (1 + F'x)D^4 + 2(H + H'u)R^2D^2, \quad (25)$$

where scaled parameters are defined by

$$G = -\frac{g_R}{\sqrt{g'_R}}, \quad F = -\frac{g_D}{\sqrt{g'_D}}, \quad F' = -\frac{h_D}{g'_D}, \quad H = \frac{g''_D}{2\sqrt{g'_D g'_R}}, \quad H' = \frac{h''_D}{2\sqrt{g'_D g'_R}}. \quad (26)$$

For large values of R^2 and D^2 , eq. (25) behaves as a linear quadratic form of them for any fixed values of (x, u) : $V_{\text{eff}} \sim R^4 + (1 + F'x)D^4 + 2(H + H'u)R^2D^2$. The Sylvester's theorem should give the positive definite condition for it [12]:

$$\text{a) } 1 + F'x - (H + H'u)^2 \geq 0$$

or

$$\text{b) } 1 + F'x \geq 0 \quad \text{and} \quad H + H'u \geq 0, \quad (27)$$

for any values of (x, u) , the domain of which have been shown in figure 1(a) and (b). The second condition b) in eq. (27) should be required because $R^2, D^2 \geq 0$.

The condition (27) gives constraints for the possible values of parameters F' , H and H' , which can be obtained analytically. We sketch the derivation of them in Appendix C, and only show the results here.

Stability conditions in (H, F') space for fixed c and H' are classified at first into two types for $(0 \leq c \leq 2)$ and $(2 \leq c)$, and then into four types by the values of H' in each of the first two (which correspond to four subscripts I~IV in figure 2 and figure 3), which are shown in figure 2(a)~(d) for $0 \leq c \leq 2$, in figure 3(a)~(d) for $2 \leq c$. In the following lists, we obtain complete stability conditions for the coefficients F' , H , H' and c in eq. (25).

A. $0 \leq c \leq 2$ case

$$\text{a : } H' \leq -\frac{3}{2-c} \quad (\text{figure 2(a)})$$

$$F' \geq -3, \quad (-H' - 1 \leq H) \quad (28)$$

This critical point $(-H' - 1, -3)$ corresponds to the point A in figure 2(a).

$$\text{b : } -\frac{3}{2-c} \leq H' \leq 0 \quad (\text{figure 2(b)})$$

$$F' \geq k(c), \quad \left(-H' - 1 \leq H \leq -\frac{1+c}{3}H' \right) \quad \overline{AB} \quad (29)$$

$$F' \geq -3, \quad \left(-\frac{1+c}{3}H' \leq H \right) \quad \overline{B\infty} \quad (30)$$

$$\text{where } k(c) = \frac{3}{2} \left[(H + H') \left(H + \frac{2c-1}{3}H' \right) - 1 - \sqrt{\{(H + H')^2 - 1\} \left\{ \left(H + \frac{2c-1}{3}H' \right)^2 - 1 \right\}} \right].$$

$$\text{c: } 0 \leq H' \leq \frac{3}{2+c} \quad (\text{figure 2(c)})$$

$$F' \geq \frac{(cH')^2}{1-H^2}, \quad (-1 \leq H \leq H_1) \quad \overline{\infty A} \quad (31)$$

$$F' \geq (2H - cH')^2 - 4, \quad (H_1 \leq H \leq H_2(c)) \quad \overline{AB} \quad (32)$$

$$F' \geq k(-c), \quad \left(H_2(c) \leq H \leq \frac{c-1}{3}H' \right) \quad \overline{BC} \quad (33)$$

$$F' \geq -3, \quad \left(\frac{c-1}{3}H' \leq H \right) \quad \overline{C\infty} \quad (34)$$

$$\text{where } H_1 \equiv \frac{cH' - \sqrt{16+c^2H'^2}}{4}, \quad H_2(c) = \frac{(3c+2)H' - \sqrt{16+(c+2)^2H'^2}}{4}.$$

d: $\frac{3}{2+c} \leq H'$ (figure 2(d))

$$F' \geq \frac{(cH')^2}{1-H^2}, \quad (-1 \leq H \leq H_1) \quad \overline{\infty A} \quad (35)$$

$$F' \geq (2H - cH')^2 - 4, \quad \left(H_1 \leq H \leq \frac{cH' - 1}{2} \right) \quad \overline{AB} \quad (36)$$

$$F' \geq -3, \quad \left(\frac{cH' - 1}{2} \leq H \right) \quad \overline{B\infty} \quad (37)$$

B. $2 \leq c$ case

a: $H' \leq -\frac{3}{c-2}$ (figure 3(a))

$$F' \geq (2H + cH')^2 - 4, \quad \left(-H' - 1 \leq H \leq -\frac{1+cH'}{2} \right) \quad \overline{AB} \quad (38)$$

$$F' \geq -3, \quad \left(-\frac{1+cH'}{2} \leq H \right) \quad \overline{B\infty} \quad (39)$$

b: $-\frac{3}{c-2} \leq H' \leq 0$ (figure 3(b))

$$F' \geq (2H + cH')^2 - 4, \quad (-H' - 1 \leq H \leq H_2(-c)) \quad \overline{AB} \quad (40)$$

$$F' \geq k(c), \quad \left(H_2(-c) \leq H \leq -\frac{1+c}{3}H' \right) \quad \overline{BC} \quad (41)$$

$$F' \geq -3, \quad \left(-\frac{1+c}{3}H' \leq H \right) \quad \overline{C\infty} \quad (42)$$

c: $0 \leq H' \leq \frac{3}{2+c}$ (figure 3(c))

$$F' \geq \frac{(cH')^2}{1-H^2}, \quad (-1 \leq H \leq H_1) \quad \overline{\infty A} \quad (43)$$

$$F' \geq (2H - cH')^2 - 4, \quad (H_1 \leq H \leq H_2(c)) \quad \overline{AB} \quad (44)$$

$$F' \geq k(-c), \quad \left(H_2(c) \leq H \leq \frac{c-1}{3}H' \right) \quad \overline{BC} \quad (45)$$

$$F' \geq -3, \quad \left(\frac{c-1}{3}H' \leq H \right) \quad \overline{C\infty} \quad (46)$$

d: $\frac{3}{2+c} \leq H'$ (figure 3(d))

$$F' \geq \frac{(cH')^2}{1-H^2}, \quad (-1 \leq H \leq H_1) \quad \overline{\infty A} \quad (47)$$

$$F' \geq (2H - cH')^2 - 4, \quad \left(H_1 \leq H \leq \frac{cH' - 1}{2} \right) \quad \overline{AB} \quad (48)$$

$$F' \geq -3, \quad \left(\frac{cH' - 1}{2} \leq H \right) \quad \overline{B\infty} \quad (49)$$

VI. APPLICATIONS

In this section we show some applications to our model. As a first application, we obtain the phase diagrams for the two special sets of the coefficients (26). As a second application, we construct a kinetic term by introducing a coupling to the gauge boson and study the Higgs phenomenon in the three characteristic phases.

A. Phase diagrams for some cases

1. The angular parameter (x, u) independent potential

We study the condensed phases in the case of (x, u) -independent potential, $F' = 0$ and $H' = 0$ in eq. (25). The effective potential becomes a function of R and D with three parameters (G, F, H) :

$$V(R, D) = -GR^2 + R^4 - FD^2 + D^4 + 2HR^2D^2. \quad (50)$$

The stability conditions are given as in the previous section:

$$1 - H^2 \geq 0 \quad \text{or} \quad H \geq 0. \quad (51)$$

Four kinds of condensed phase (as extremum points of the effective potential) are obtained: 1) $R_0 = D_0 = 0$, 2) $R_0 \neq 0, D_0 = 0$, 3) $R_0 = 0, D_0 \neq 0$, 4) $R_0 \neq 0, D_0 \neq 0$ corresponding to eqs. (4)-(7), and the parameter regions for each phase are shown in Table 1 with the conditions (51). Referring to Table 1, the phase diagram on (F, G) space for fixed H is obtained in figure 4 by comparing $V(R_0, D_0)$ in the ranges where more than two kinds of extrema exist. At $H=-1$, the mixed phase emerges above the line $\overline{AB}$ ($G=-F$) in figure 4(a), and as H increases to 1 the two branches of this line merges into the line $\overline{OA}$ ($G = F$) in figure 4(c). When H exceeds 1 the mixed phase vanishes.

2. The potential for the symmetric phase

Next we study the case in which only the symmetric **6** phase emerges, i.e., $g_R=g'_R=g''_D=h''_D=0$ in eq. (12). In this case the effective potential is given by

$$V(x, D) = g_D D^2 + (g'_D - h_D x) D^4. \quad (52)$$

From $0 \leq x \leq \frac{1}{3}$, we obtained the stability condition by

$$h_D \leq 3 g'_D, \quad g'_D \geq 0. \quad (53)$$

This condition coincides with the previous results in $F \geq -3$ and $H'=H=0$, e.g., in eqs. (29) and (30). From eq. (52) the four kinds of condensed phase are obtained: 1) $D_0=0, x=0$, 2) $D_0 \neq 0, x=0$, 3) $D_0 \neq 0, x=\frac{1}{3}$. The parameter regions and the form of the order parameter S_0 in eq. (9) for each phase are given in Table 2. The phase diagram in (F, G) space for $g_D \geq 0$ is shown in figure 5.

It is noticed that the linear dependence of the effective potential (52) on x is the reason why the phase characterized by, e.g., $S_0=(d_1, d_2, 0)$ does not emerge.

B. Higgs phenomenon

Higgs phenomenon is the one where gauge bosons acquire finite effective masses in the broken-symmetry phase. In the minimal coupling scheme, the gauge boson fields A_μ^a are introduced through the kinetic terms of the order parameter. To the second-order, it is given by

$$K_{kinetic} = -k_A \text{Tr}(D_\mu A)^* D^\mu A + k_S \text{Tr}(D_\mu S)^* D^\mu S, \quad (54)$$

where the coefficients k_A and k_S are positive constants. The covariant derivative D_μ in eq. (54) is defined by

$$D_\mu \Psi \equiv \partial_\mu \Psi - ig \frac{\lambda_a}{2} A_\mu^a \Psi - ig \Psi \frac{t \lambda_a}{2} A_\mu^a, \quad (55)$$

where g and $\lambda_{a=1\sim 8}$ are the gauge coupling constant and the Gell-Mann matrices [14].

The effective masses of the gauge bosons are obtained from the second-order terms of the gauge boson fields in the kinetic terms (54).

1. The $\mathbf{3}^*$ condensed phase

In the case of the $\mathbf{3}^*$ condensed phase we consider an order parameter

$$\Psi_0 = \begin{pmatrix} 0 & -a & 0 \\ a & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (56)$$

where a is a real constant. Note that this form is general enough since other forms of the $\mathbf{3}^*$ phase is obtained by a simple transformation to Ψ_0 as in eq. (8). In this phase $SU_c(3)$ symmetry breaks into $SU(2)$.

By substituting Ψ_0 into eq. (54) the mass terms are obtained: $m_v^2 \text{Tr} v^\dagger v$ and $m_\phi^2 \text{Tr} \phi^2$ where $m_v \equiv \sqrt{2k_A} |a| g$, $m_\phi \equiv \sqrt{\frac{4}{3}k_A} |a| g$, $v = \frac{1}{2}(\lambda_4 + i\lambda_5)(A_\mu^4 + iA_\mu^5) + \frac{1}{2}(\lambda_6 + i\lambda_7)(A_\mu^6 + iA_\mu^7)$ and $\phi = \lambda_8 A_\mu^8$. The gauge fields v and ϕ are different multiplets of the residual symmetry $SU(2)$.

This phase is the one which is featured in most microscopic models. Using the values of the gauge coupling constant g and the gap $|a|$ calculated in models [4], the effective masses are estimated: $m_v = \sqrt{\frac{3}{2}}m_\phi \sim 100$ MeV.

2. The **6** condensed phase

We select an order parameter proportional to the identity matrix which is discussed in the previous application, see Table 2 :

$$\Psi_0 = \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix}. \quad (57)$$

In this phase $SU(3)$ symmetry breaks into $SO(3)$. Since the gauge boson field $A_\mu (\equiv \lambda_a A_\mu^a)$ is transformed as direct products of two triplets under the $SO(3)$ group, they are decomposed into the direct sum of triplet (anti-symmetric part) and quintet (symmetric part):

$$A_\mu = V_{\mathbf{3}} + T_{\mathbf{5}}, \quad (V_{\mathbf{3}} \equiv \sum_{a=2,5,7} \lambda_a A_\mu^a), \quad (T_{\mathbf{5}} \equiv \sum_{a=1,3,4,6,8} \lambda_a A_\mu^a). \quad (58)$$

Since $T_{\mathbf{5}}$ is assigned to the generators of the broken symmetry $SU(3)/SO(3)$, the mass term is accompanied with the quintet term: $m_T^2 \text{Tr}(T_{\mathbf{5}})^2$, where $m_T \equiv \sqrt{4k_S}|a|g$.

3. The Mixed condensed phase

We study a special case for the mixed phase characterized by the order parameter:

$$\Psi_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ a & 0 & 0 \end{pmatrix}. \quad (59)$$

In this phase $SU(3)$ symmetry breaks into $U(1)$ which is generated by $H(t) = e^{i(\lambda_3 + \sqrt{3}\lambda_8)t}$. Since in this case the multiplets of the gauge field are not trivial, we investigate a dependence of the gauge field A_μ on the infinitesimal transformation of $H(t)$.

$$(HA_\mu H^\dagger)_{kl} \sim (A_\mu)_{kl} + (P)_{kl}(A_\mu)_{kl}, \quad P = i \begin{pmatrix} 0 & 1 & 2 \\ -1 & 0 & 1 \\ -2 & -1 & 0 \end{pmatrix} t \quad (60)$$

Noticing that $(\lambda_3 + \sqrt{3}\lambda_8)$ and $(\lambda_3 - \frac{1}{\sqrt{3}}\lambda_8)$ are orthogonal each other, the gauge field are decomposed into the multiplets:

$$A_\mu = \begin{pmatrix} 0 & \xi & \eta \\ \xi^* & 0 & \zeta \\ \eta^* & \zeta^* & 0 \end{pmatrix} + (\lambda_3 - \frac{1}{\sqrt{3}}\lambda_8)\varphi + (\lambda_3 + \sqrt{3}\lambda_8)\varphi', \quad (61)$$

$$\xi \equiv A_\mu^1 - iA_\mu^2, \quad \eta \equiv A_\mu^4 - iA_\mu^5, \quad \zeta \equiv A_\mu^6 - iA_\mu^7$$

$$\varphi \equiv (3A_\mu^3 - \sqrt{3}A_\mu^8)/4, \quad \varphi' \equiv (A_\mu^3 + \sqrt{3}A_\mu^8)/4.$$

The mass terms are summerized as follows:

$$m_\xi^2 \xi^* \xi = \frac{1}{2}(k_A + k_S)|a|^2 g^2 \xi^* \xi, \quad m_\eta^2 \eta^* \eta = 2k_S|a|^2 g^2 \eta^* \eta$$

$$m_\zeta^2 \zeta^* \zeta = \frac{1}{2}(k_A + k_S)|a|^2 g^2 \zeta^* \zeta, \quad m_\varphi \varphi^2 = \frac{16}{18}(k_A + k_S)|a|^2 g^2 \varphi^2. \quad (62)$$

In the same way more complicated cases can also be studied.

VII. SUMMARY

We have derived the general form of the Color- $SU(3)$ -Ginzburg-Landau Effective Potential for the order parameter with $\mathbf{3} \times \mathbf{3}$ symmetry to the fourth-order and determined the range of coefficients which stabilizes the vacuum of the system. Corresponding to the irreducible components of the order parameter $\mathbf{3} \times \mathbf{3} = \mathbf{3}^* + \mathbf{6}$, we found four kinds of phases: non condensed phase, anti-symmetric ($\mathbf{3}^*$) phase, symmetric ($\mathbf{6}$) phase and mixed ($\mathbf{3}^* + \mathbf{6}$) phase.

As applications, we obtained the phase diagrams in the two special cases and studied the Higgs phenomenon in some characteristic phases by gauging this model.

The full classification of the broken vacuum are analytically possible including the angular parameter x and u , and will be discussed in different paper.

APPENDIX A: NORMAL COORDINATE DECOMPOSITION

In this appendix, we show the decomposition of the 3-by-3 symmetric matrix $S = {}^tS$ in eq. (9). The proof is a variation of that used in the group representation theory [13], the $SU(3)$ -chiral nonlinear sigma model [14] and the Kobayashi-Maskawa theory of quark mass matrix [15].

We take the subsidiary hermite matrix $H = S^*S = S^\dagger S$. Its eigenvalue λ_i and eigenvectors $\mathbf{e}_i$ are obtained from the eigenequation: $H\mathbf{e}_i = \lambda_i\mathbf{e}_i$ ($i = 1, 2, 3$).

In this appendix, we consider only the case of no degeneracy: $\lambda_i \neq \lambda_j$ ($i \neq j$). (The extension to degenerate cases should be made by analytic continuation.)

Because of the hermiticity of H , the eigenvectors $\mathbf{e}_i$ can be taken to be orthogonal unit vectors (${}^\dagger\mathbf{e}_i\mathbf{e}_j = \delta_{ij}$), and the eigenvalues λ_i take real positive values. (because of $H = S^\dagger S$).

In matrix form, the eigenequation becomes

$$U^\dagger H U = L \equiv \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}, \quad (\text{A1})$$

where $U = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3) \in SU(3)$.

The eigenvector $\mathbf{e}_i$ satisfies

$$H^* S \mathbf{e}_i = S S^* S \mathbf{e}_i = S H \mathbf{e}_i = \lambda_i S \mathbf{e}_i. \quad (\text{A2})$$

so that we obtain $S \mathbf{e}_i = \alpha_i \mathbf{e}_i^*$ (because $H^* \mathbf{e}_i^* = \lambda_i \mathbf{e}_i^*$). $\alpha_i \equiv |\alpha_i| e^{i\theta_i}$ are complex numbers. In matrix form, it becomes

$${}^t U S U = Z \equiv \begin{pmatrix} \alpha_1 & 0 & 0 \\ 0 & \alpha_2 & 0 \\ 0 & 0 & \alpha_3 \end{pmatrix}. \quad (\text{A3})$$

The polar decomposition of Z becomes

$$A = R e^{iT} = \begin{pmatrix} |\alpha_1| & 0 & 0 \\ 0 & |\alpha_2| & 0 \\ 0 & 0 & |\alpha_3| \end{pmatrix} \begin{pmatrix} e^{i\theta_1} & 0 & 0 \\ 0 & e^{i\theta_2} & 0 \\ 0 & 0 & e^{i\theta_3} \end{pmatrix}, \quad T = \begin{pmatrix} \theta_1 & 0 & 0 \\ 0 & \theta_2 & 0 \\ 0 & 0 & \theta_3 \end{pmatrix}. \quad (\text{A4})$$

Because of $\lambda_i \mathbf{e}_i = H \mathbf{e}_i = S^* S \mathbf{e}_i = \alpha_i S^* \mathbf{e}_i^* = \alpha_i (S \mathbf{e}_i) = |\alpha_i|^2 \mathbf{e}_i$, we obtain $\lambda_i = |\alpha_i|$ and also $R = \sqrt{L}$. The polar decomposition of A becomes $A = \sqrt{L} e^{iT}$.

Because the angular part T is diagonal, it can be expanded by the 3-by-3 unit matrix I and the diagonal Gell-Mann matrices $\lambda_{3,8}$: $T = \theta I + \phi_3 \lambda_3 + \phi_8 \lambda_8$. If we define $K = e^{\frac{i}{2}(\phi_3 \lambda_3 + \phi_8 \lambda_8)}$, then we obtain $e^{iT} = K e^{i\theta} I K$. Using the first equation in (A4), the matrix S becomes

$$S = U^* Z U = U^* \sqrt{L} e^{iT} U^\dagger = e^{i\theta} U^* K \sqrt{L} K U^\dagger = e^{i\theta} G \sqrt{L} {}^t G, \quad (\text{A5})$$

where $G \equiv U^* K$ and ${}^t G = {}^t K U^\dagger = K U^\dagger$. This is just the decomposition of S in eq. (8) with $S_0 = \sqrt{L}$.

If we define the matrix A_0 by $A = G A_0 {}^t G$, then A_0 also becomes antisymmetric and is generally represented as the first equation in eq. (9).

APPENDIX B: MAXIMUM AND MINIMUM OF f

Instead of p_i with $p_1 + p_2 + p_3 = 1$, we take independent variables (s, t) :

$$s = \frac{1}{\sqrt{2}}(-p_1 + p_2), \quad t = -\sqrt{\frac{3}{2}} \left(p_1 + p_2 - \frac{2}{3} \right). \quad (\text{B1})$$

The possible range of (s, t) is determined from the condition: $0 \leq p_i \leq 1$ and $p_1 + p_2 + p_3 = 1$. Using (B1), it is found to be the inside of the equilateral triangle with vertices $(0, \sqrt{2/3})$ and $(\pm 1/\sqrt{2}, -1/\sqrt{6})$ in the (s, t) plane. Since the function $f(s, t)$ is an even function of s , $f(s, t) = f(-s, t)$, we may consider a half ($s \geq 0$) of the triangular region (figure 6) to find its maximum/minimum.

For a fixed value of x , a trajectory of $x = p_2 p_3 + p_3 p_1 + p_1 p_2 = -(s^2 + t^2)/2 + 1/3$ in (s, t) -plane become a circle with the center $(0, 0)$ and the radius $\sqrt{(2/3) - 2x}$ (figure 6). From the aspect of intersections between the circle and the triangular region, two cases should be discriminated:

1. When $0 \leq x < 1/4$, the circle intersects with three sides of the triangle (figure 6(a)), and the two separate arcs (B_0B_1 and B_2B_3 in figure 6(a)) are included in the half triangular region.

2. When $1/4 \leq x \leq 1/3$, the whole circle is included in the triangular region (figure 6(b)).

The intersections with the s -axis ($t = 0$) are denoted by B_0 and B_4 .

The (s, t) -coordinates of the above four points $B_{0\sim 4}$ are given by

$$B_0 = \left(0, \sqrt{\frac{2}{3} - 2x}\right), \quad B_1 = \left(\frac{1 - \sqrt{1 - 4x}}{2\sqrt{2}}, \frac{1 + 3\sqrt{1 - 4x}}{2\sqrt{6}}\right), \quad (\text{B2})$$

$$B_2 = \left(\frac{1 + \sqrt{1 - 4x}}{2\sqrt{2}}, \frac{1 - 3\sqrt{1 - 4x}}{2\sqrt{6}}\right), \quad B_3 = \left(\sqrt{\frac{1 - 4x}{2}}, -\frac{1}{\sqrt{6}}\right), \quad (\text{B3})$$

$$B_4 = \left(0, -\sqrt{\frac{2}{3} - 2x}\right). \quad (\text{B4})$$

Let's consider the maximum value of f_3^+ for fixed values of c and x . Using $s = \sqrt{6 - 9t^2 - 18x}/3$ and (B4), f_3^+ in eq. (19) becomes

$$f_3^+(t) = \frac{1}{3} \left[\sqrt{6}t + 1 + c\sqrt{6t^2 - \sqrt{6}t + 9x - 2} \right], \quad (\text{B5})$$

where we take t as an independent variable and (x, c) as any fixed parameters. Solving the equation $[f_3^+(t)]' = 0$, we obtain one local extremum at $B_M = (\sqrt{6 - 9t_M^2 - 18x}/3, t_M)$ where

$$t_M = \frac{1}{2\sqrt{6}} \left[1 - 3\sqrt{\frac{(4x - 1)(c^2 - 1)}{1 - c^2}} \right]. \quad (\text{B6})$$

for some values of (x, c) . The parameter range where the extremum of f_3^+ exists (at B_M) can be read off from the condition that t_M in (B6) should take a real value: i.e., $(4x - 1)(c^2 - 1) \geq 0$. Solving this inequality, we obtain two separated regions: a) $c \leq 1$ & $x \leq 1/4$ and b) $c \geq 1$ & $x \geq 1/4$ (Shaded regions in figure 7) where the function f_3^+ has an extremum. Judging from the signature of $[f_3^+]''(t_M)$, we can find that $f_3^+(t_M)$ is a local maximum in region a) and minimum in region b). For t_M have to be in the permissible region shown in figures 6(a) and (b), we obtain further condition:

1. When $x \leq 1/4$ and $c \leq 1$, t_M have to be on the arc B_2B_3 (figure 6(a)). Thus, we obtain $x \geq c^2/4$ ($\overline{OD}$ in figure 7).
2. When $x > 1/4$, t_M have to be on the half circle B_0B_4 (figure 6(b)). Thus, we obtain $c \leq 2$ or $x \leq c(5c-4)/[4(2c-1)^2]$ ($\overline{AB}$ in figure 7),

Summarizing the above discussions about the aspects of extrema of f_3^+ , we can classify the five regions in the (c, x) -plane (figure 7).

Let's analyze maximum points in each regions. Candidates of the absolute maximum point of f_+^3 are the end points $B_{0\sim 4}$ where

$$\begin{aligned}
f_+^3(B_0) &= \frac{1+c}{3} + \frac{2-c}{3}\sqrt{1-3x}, & f_+^3(B_1) &= \frac{1}{2} + \frac{1}{2}\sqrt{1-4x}, \\
f_+^3(B_2) &= \frac{1}{2} - \frac{1}{2}\sqrt{1-4x}, & f_+^3(B_3) &= c\sqrt{x}, \\
f_+^3(B_4) &= \frac{1+c}{3} - \frac{2-c}{3}\sqrt{1-3x},
\end{aligned} \tag{B7}$$

or the local maximum at M (if exist):

$$f_+^3(B_M) = \frac{1}{2} - \frac{1}{2}\sqrt{(c^2-1)(4x-1)}. \tag{B8}$$

Comparing $f_+^3(B_i)$ in eqs. (B7) and (B8), we can find maximum points in each region in figure 7:

1. The maximum is at B_0 in regions 1, 2, a part of 3 ($c \leq 2$ or $x \leq (2c-1)^2/[4(1-c+c^2)^2]$ $\overline{EF}$ in figure 8), and a part of 4 ($c \leq 2$).
2. The maximum is at B_3 in a remaining part of region 3.
3. The maximum is at B_4 in region 5 and a remaining part of 4.

The results are summarized in figure 8.

In a similar way, the minimum value of $f_3^-(t)$ can be calculated:

$$f_3^-(t) = \begin{cases} c\sqrt{x} & (0 \leq x < 1/4) \\ \frac{1-c}{3} - \frac{2+c}{9}\sqrt{1-3x} & (1/4 \leq x \leq 1/3) \end{cases} \tag{B9}$$

APPENDIX C: DERIVATION OF STABILITY CONDITIONS

In this appendix, we sketch the derivation of the stability condition: a parameter range of (F', H, H') that satisfies (27).

Instead of (x, u) , we take new variables (X, U) defined by

$$X \equiv 1 + F'x, \quad U \equiv H + H'u. \quad (\text{C1})$$

Using them, the conditions in (27) become simple:

$$\text{a) } X \geq U^2, \quad \text{or} \quad \text{b) } X \geq 0 \text{ and } U \geq 0. \quad (\text{C2})$$

In the (X, U) -plane, the range that satisfies these conditions is represented by the hatched area in figure 9. We denote it by the region Σ .

The possible range of variables (X, U) , which is translated from the one of (x, u) (given in figure 1), depends on parameters (F', H', c) and we denote it by $\Lambda(F', H', c)$. In figure 9, as an example, we show the $\Lambda(F', H', c)$ as the cross-hatched region for some values of (F', H', c) that satisfies $F' \geq 0$, $H' \geq 0$ and $c \geq 2$. In (X, U) -plane, the stability condition (27) become equivalent with the geometrical condition:

$$\Lambda(F', H', c) \subset \Sigma. \quad (\text{C3})$$

The boundary of $\Lambda(F', H', c)$ and Σ are patchworks of line or parabolic segments, so that the condition (C3) can be solved algebraically.

As an example, let's consider the case that $c \leq 2$, $F' \geq 0$ and $H' \geq 0$ (figure 9). In this case, eq. (C3) can be attributed into whether the boundary segment $\overline{\infty GO}$ for Σ intersects that of Λ , $\overline{AB}$, or not. Both segments, $\overline{\infty GO}$ and $\overline{AB}$, have parabolic shapes represented by $X = U^2$ ($\overline{\infty GO}$) and $X = [F'/(cH')^2](U - H)^2 + 1$ ($\overline{AB}$). It can be judged from the condition that the edge points, $A = (1, H)$ and $B = (\frac{F'}{4} + 1, H - \frac{cH'}{2})$, are included in Σ ; it gives two necessary conditions for parameters:

$$-1 \leq H, \quad \left(H - \frac{cH'}{2} \right)^2 \leq \frac{F'}{4} + 1. \quad (\text{C4})$$

In case that $F'/(cH')^2 \leq 1$, the slope of $\overline{\infty GO}$ is more gradual than that of AB , and eqs. (C4) are sufficient condition. However, in the case that $F'/(cH')^2 \geq 1$, an extra condition should be added in order that two parabolic segments have no intersections:

$$(cH')^2 + F'(H^2 - 1) < 0. \quad (\text{C5})$$

Eqs. (C4) and (C5) give the necessary and sufficient stability condition in that case:

$$F' \geq \frac{(cH')^2}{1 - H^2}, \quad F' \geq (2H - cH')^2 - 4, \quad -1 \leq H, \quad (\text{C6})$$

which is shown as the hatched region in figure 2.

Other cases can be treated in a similar way.

REFERENCES

- [1] Barrois F 1977 *Nucl. Phys.* **B129** 390
- [2] Bailin D and Love A 1984 *Phys. Rep.* **107** 325
- [3] Iwasaki M 1996 *Prog. Theor. Phys.* **96** 1043
- [4] Rapp R, Schäfer T, Shuryak E and Velkovsky M 1998 *Phys. Rev. Lett.* **81** 53
 Alford M, Rajagopal K and Wilczek F 1998 *Phys. Lett.* **B422** 247.
- [5] Son D T 1999 *Phys. Rev.* **D59** 094019
 Schäfer T and Wilczek F 1998 *Phys. Lett.* **B450** 325
 Hsu S and Schwetz M 2000 *Nucl. Phys.* **B572** 211
- [6] Schäfer T and Wilczek F 1999 *Phys. Rev.* **D60** 114033
 Brown W E, Liu J T and Ren H 2000 *Phys. Rev.* **D62** 054016
 Pisarski R D and Rischke D H 2000 *Phys. Rev.* **D61** 074017
- [7] Alford M, Rajagopal K and Wilczek F 1999 *Nucl. Phys.* **B537** 443
 Shovkovy I A and Wijewardhana L C R 1999 *Phys. Lett.* **B470** 189
- [8] Casalbuoni R and Gatto R 1999 *Phys. Lett.* **B464** 111
 Rho M, Wirzba A and Zahed I 2000 *Phys. Lett.* **B473** 126
- [9] Berges J and Rajagopal K 1999 *Nucl. Phys.* **B538** 215
 Miransky V A, Shovkovy I A and Wijewardhana L C R 1999 *Phys. Lett.* **B468** 270
- [10] For example, Tilley D R and Tilley J 1986 *“Superfluidity and Superconductivity”, 2nd ed.* Adam Hilger Ltd
- [11] Iida K and Baym G 2001 *Phys. Rev.* **D63** 074018
 Giannakis I and Ren H 2002 *Phys. Rev.* **D65** 054017
 Sedrakian D M, Blaschke D, Shahabasyan K M and Voskresensky D N 2001 *Astrofiz.* **44** 443

- [12] Gradshteyn I S and Ryzhik I M 1990 *“Tables of Integral, Series, and Products”, 5th ed.*
(translated from Russian by Scripta Technica Inc.) San Diego, Academic Press
- [13] Murnaghan F D 1938 *“The Theory of Group Representations”* Baltimore, The John Hopkins Press
- [14] Lee B 1972 *“Chiral Dynamics”* New York, Gordon and Breach Sci. Pub.
- [15] Kobayashi M and Maskawa T 1973 *Prog. Theor. Phys.* **49** 652

figure 1. Range of variables x and u . a) for $0 \leq c \leq 2$, b) for $2 \leq c$. Details are given in the text.

figure 2. Stability region in (H, F') space for $(0 \leq c \leq 2)$ (the shaded area). figure 2(a) is for $H' \leq -\frac{3}{2-c}$, (b) for $-\frac{3}{2-c} \leq H' \leq 0$, (c) for $0 \leq H' \leq \frac{3}{2+c}$ and (d) for $\frac{3}{2+c} \leq H'$.

figure 3. Stability region in (H, F') space for $(2 \leq c)$. figure 3(a) is for $H' \leq -\frac{3}{c-2}$, (b) for $-\frac{3}{c-2} \leq H' \leq 0$, (c) for $0 \leq H' \leq \frac{3}{2+c}$ and (d) for $\frac{3}{2+c} \leq H'$.

figure 4. (a) The phase diagram for the effective potential (50) in (F, G) space when $H=-1$. $\overline{AB} : G=-F$. The regions with vertical stripes, with horizontal stripes and with cross hatch show $\mathbf{3}^*$, $\mathbf{6}$ and the Mixed condensed phases. The remaining part ($G \leq 0$ and $F \leq 0$) shows the Non condensed phase. (b) The same, but $H=1/2$. $\overline{OA} : G=2F$. $\overline{OB} : G=F/2$. (c) The same, but $H=1$. $\overline{OA} : G=F$.

figure 5. The phase diagram for the effective potential (52) in (g'_D, h_D) space when $g_D < 0$. $\overline{AB} : h_D = 3g'_D$. The regions I and II corresponds to the phases 2) and 4) in Table 2, but the region just on the line $h_D=0$ ($g'_D > 0$) corresponds to the phase 3).

figure 6. Range of variables s and t is shown as inside of the half triangle ACD. Trajectories of (s, t) for fixed x are given for $0 \leq x \leq \frac{1}{4}$ by two separate arcs B_0B_1 and B_2B_3 in (a) and for $\frac{1}{4} \leq x \leq \frac{1}{3}$ by a half circle B_0B_4 in (b).

figure 7. The five regions in the (c, x) -plane according to the aspects of t_M . t_M is defined in the shaded regions. In addition, in the region 1 and 4 t_M exists on the arc B_2B_3 (figure 6(a)), and on the half circle B_0B_4 (figure 6(b)).

figure 8. The three regions in the (c, x) -plane separated by the maximum values of f_+^3 , which are $f_+^3(B_0)$, $f_+^3(B_3)$ and $f_+^3(B_4)$ in the region I, II and III.

figure 9. An illustration for the stability condition $\Lambda(F', H', c) \subset \Sigma$ in the case $c \leq 2$, $F' \geq 0$ and $H' \geq 0$.

Table 1 : Condensed phases and their parameter ranges.

	R_0^2	D_0^2	$V(R_0, D_0)$	the ranges where R_0, D_0 are defined
Non cond.	0	0	0	$G \leq 0, \quad F \leq 0$
3* cond.	$G/2$	0	$-G^2/4$	$G \geq 0$
6 cond.	0	$F/2$	$-F^2/4$	$F \geq 0$
Mix cond.	$\frac{G-HF}{2(1-H^2)}$	$\frac{F-HG}{2(1-H^2)}$	$\frac{-G^2-F^2+2FGH}{4(1-H^2)}$	$\left\{ \begin{array}{l} G \geq HF \text{ and } G \geq F/H, \text{ if } -1 \leq H \leq 0 \\ G \geq HF \text{ and } G \leq F/H, \text{ if } 0 \leq H \leq 1 \\ G \leq HF \text{ and } G \geq F/H, \text{ if } 1 \leq H \end{array} \right.$

Table 2 : Forms of the order parameter and their parameter ranges

(D_0^2, x)	$S_0 = (d_1, d_2, d_3)$	the parameter region
1) $(0, 0)$	$d_1 = d_2 = d_3 = 0$	$g_D \geq 0$
2) $(-\frac{g_D}{2g'_D}, 0)$	$d_i \neq 0, d_{j \neq i} = 0$	$g_D < 0, h_D < 0$
3) $(-\frac{g_D}{2g'_D}, x)$	$d_1^2 + d_2^2 + d_3^2 = D_0^2$	$g_D < 0, h_D = 0$
4) $(-\frac{g_D}{2(g'_D - h_D/3)}, \frac{1}{3})$	$d_1 = d_2 = d_3 \neq 0$	$g_D < 0, 0 < h_D \leq 3g'_D$

figure 1




figure 2







figure 3




figure 4




figure 5




figure 6





figure 7




figure 8




figure 9

